



EVALUATION OF HEPATIC BIOMARKERS IN PREGNANT WOMEN WITH PREECLAMPSIA

Seyyed Hossein Hassanpour^{1*} and Seyyed Zeinab Karami²

¹Young Researchers and Elite Club, Yasooj Branch, Islamic Azad University, Yasooj, Iran

²Department of Biology, Faculty of Basic Sciences, Yasouj University, Yasouj, Iran

ARTICLE INFO

Article History:

Received 20th May, 2017

Received in revised form 23rd

June, 2017

Accepted 27th July, 2017

Published online 28th August, 2017

Key words:

Preeclampsia, HELLP, ALT,
AST, Bilirubin

ABSTRACT

Preeclampsia is a pregnancy-related disorder and considered as one of major reasons of infants and mothers death in developed nations. HELLP disorder is a problem related to child birth that usually happens in women with intense preeclampsia and associated with different features, including hemolysis, elevated liver enzymes and low platelet count. Due to normal hepatic markers during pregnancy, our purpose is to examine these factors in pregnant women and their association with disorders such as preeclampsia and HELLP syndrome. This case-control study included 99 Iranian pregnant women that divided two group including preeclamptic and normotensive pregnant women. Samples were collected from Ahvaz city. We measured liver enzymes activity (ALT, AST and ALP), total bilirubin and direct bilirubin and blood platelets by calorimetric method in both groups. The results showed that there was no significant difference in the platelet level in both groups. However we found significant difference in the serum level of ALT, AST, ALP and total bilirubin between two groups ($p < 0.05$), while the result related to direct bilirubin was not significant at the end of study. The outcomes related to this study show that hepatic biomarkers in pregnant women with preeclampsia higher than normal pregnant women therefore, we can predict more likely to develop HELLP syndrome in pregnant women with preeclampsia.

Copyright © 2017 Seyyed Hossein Hassanpour and Seyyed Zeinab Karami. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Preeclampsia is a pregnancy-related disorder and considered as one of major reasons of infants and mothers death in developed nations.^[1] Some of risk factors for development of preeclampsia are diabetes mellitus, hypertension, obesity and anti-phospholipid antibody syndrome.^[2] Each year, 585,000 women die due to complications related to pregnancy that is 95% of them in developing countries and among them, 30% of cases are due to problems of hypertension during pregnancy and particularly preeclampsia.^[3] In the USA, 18% of mortality among women is related to pregnancy period and especially hypertension disorders during this period.^[4] Generally, the definition of preeclampsia is complicated due to differences in its diagnosis, but persistence in blood pressure that occurs in 12-22% of pregnancies and is dependent on the type of population can be one of the its reasons.^[5] Also vascular endothelial dysfunction is final common pathway that causes the mother's system response.^[1] For the example, one study was showed that in women with preeclampsia, increase in level of soluble fms-like tyrosine kinase 1 (sFlt-1), and a decrease in PlGF levels is higher than the control group, as these changes occur several weeks before the first onset, their measurement can be a good predictors.^[6] Researchers has been found that men and women, whose mothers during pregnancy

have preeclampsia, likely their children develop preeclampsia later that shows the effect of genetic factors in this disorder, although not too much information in this field.^[7] In addition, in a study was conducted in Iran, winter and urinary tract infection were considered as risk factors for preeclampsia.^[8] It has been reported that slightly changes occur in biomarkers of liver during pregnancy, indeed in this period, the level of AST, ALT, GGT, serum bilirubin and bile acids usually remain within normal range, therefore any change in their level may indicate a problem.^[9] HELLP disorder is a problem related to child birth that usually happens in women with intense preeclampsia.^[10] This syndrome occurs mainly in preterm and sometimes in during late gestation and after child birth.^[11] HELLP is expressed as the following three features: hemolysis, elevated liver enzymes and low platelet count.^[12] This syndrome is a very dangerous situation and leads to serious problems such as hemolysis, epigastric pain, decrease of liver enzymes and thrombocytopenia in during this syndrome.^[10] In Iran, a study conducted on preeclampsia was showed that there is a relationship between preeclampsia and vitamin D.^[13] In addition, Shahbazian *et al*, 2014 examined the relationship between preeclampsia and hypertension and microalbuminuria.^[14] As the women health is important during pregnancy and there are few studies on pregnancy and liver problems in Iran, we studied liver markers in pregnant women

and their association with disorders such as preeclampsia and HELLP syndrome. In addition, this study was in line with our recent studies on women health, including cell cycle arrest in ovarian cancer,^[15] effect of purslane extract on antioxidant balance in women with type 2 diabetes,^[16] changes in level of AGEs and β 2-microglobulin and imbalance of trace elements in type 2 diabetes.^[17,18]

MATERIAL AND METHODS

Subjects

This case-control study was performed on 99 Iranian pregnant females, whom divided two group including preeclampsia and normotensive pregnant women.

Measurements

We measured enzyme activity of liver biomarkers (ALT, AST and ALP), total bilirubin and direct bilirubin by calorimetric method and blood platelets by Hematology Analyzer - Sysmex KX-21 in pregnant women with preeclampsia (n=50) and normotensive (n=49). HELLP syndrome among woman with preeclampsia by the following criteria: total bilirubin \geq 0.6 mg/dl for detection of hemolysis, AST \geq 20 IU /L, ALT \geq 15 IU /L for the diagnosis of liver damage and blood platelet count less than 50,000 cells/ μ L.

Statistical analysis

The data were expressed as mean \pm standard deviation. For comparison of groups was used independent t test and Mann-Whitney test for platelet count and serum ALT, AST, ALP, total bilirubin, direct bilirubin, respectively. The different level was set as P<0.05.

RESULTS

The outcomes of present study was reported for 99 pregnant women including preeclampsia (n=50) and normotensive (n=49) case. The summary of these results is presented in table 1.

Table1 Comparison factors related to preeclampsia in both groups.

Factor	Group	
	Preeclampsia (n=50)	Normotensive (n=49)
Blood platelet (cell/ μ L)	223.80 \pm 72.63	216.45 \pm 47.48
ALT (IU/L)	34.34 \pm 12.77*	14.51 \pm 3.93
AST (IU/L)	41.10 \pm 10.61*	20.55 \pm 6.82
ALP (IU/L)	397.20 \pm 174.49*	180.02 \pm 46.72
Direct bilirubin (mg/dl)	0.23 \pm 0.08	0.15 \pm 0.4
Total bilirubin (mg/dl)	0.99 \pm 0.91*	0.42 \pm 0.13

Values are expressed as mean \pm SD; comparisons were made using independent t test and Mann-Whitney test for platelet count and serum ALT, AST, ALP, total bilirubin, direct bilirubin, respectively.

* Significant different with Normotensive group (P < 0.05).

The current data show that there was no significant difference in the platelet levels between normotensive pregnant and preeclampsia pregnant women. However, we obtained significant difference in the ALT serum level between normotensive pregnant women and preeclamptic pregnant women at the end of study (P<0.05) also it was found a significant difference in the AST level between two groups (P<0.05). The evaluation of serum ALP serum level was also indicated that its level in preeclamptic pregnant women was significantly higher than normotensive pregnant women (about 2-fold) (P<0.05). In relation to bilirubin level (either direct or total), the result was confirmed that direct bilirubin level in

preeclamptic pregnant women had not obvious different compared to normotensive pregnant women, while the level of total bilirubin in preeclamptic pregnant women was higher than normotensive pregnant women so that it was significant (P<0.05).

DISCUSSION

Liver function tests are abnormal in 20% to 30% of pregnancies that are associated with preeclampsia,^[19,20] and are related poor motherhood and embryonic result.^[21,22] Preeclampsia is a disorder with three features: proteinuria, hypertension and edema that occur during last trimester of 5%-10% of pregnancies. Although liver problem is infrequent in this disorder, nevertheless intense preeclampsia is related to perinatal illness and death. In fact, it is the most common reason of hepatic sensitivity and liver impairment in gestation period and 2%-12% of cases will suffer of HELLP syndrome that this syndrome is expressed as the following three features: hemolysis, elevated liver enzymes, and low platelet count. liver involvement of preeclampsias required no specific therapy, although the involvement is an indicator to prevent more serious disorders such as eclampsia, hepatic rupture, or necrosis.^[22]

We studied hepatic markers and liver damage in pregnant women in the both normotensive and preeclamptic group. In agreement with studies conducted by Weinstein *et al*, 1982^[12] and Shukla *et al*, 1978^[23] our data showed that there was significant difference in the serum ALT level between normotensive pregnant women and preeclampsia pregnant women (P<0.05). In addition, line with Cerutti *et al*, 1976,^[24] Weinstein *et al*, 1982^[12] and Shukla *et al*, 1978^[23] our data indicated that there was noticeable difference in the AST level between two groups (P<0.05). Surprisingly, the current data showed that there was no significant difference in the platelet level between normotensive pregnant and preeclampsia pregnant women. But there were significant difference in the serum ALP level and total bilirubin level between the two groups (P<0.05). The various investigations have been examined the evaluation of liver function tests and liver damage in pregnant women with preeclampsia and normal pregnant women so that obtained different results. For example; in a study by Girling *et al*, 1997 stated that the rate of liver function tests are less in normal gestation than the scope of reference presently used.^[25] The results of our project also revealed that more hepatic markers such as total bilirubin, ALT and ALP in pregnant women with preeclampsia were higher than normal pregnant women. In other study on the HELLP syndrome was indicated that AST, ALT and bilirubin were abnormal.^[12] We as well as found that total bilirubin and ALT level were more in pregnant women with preeclampsia, but direct bilirubin level had not significant difference compared to normal group. It has been reported AST level abnormality more than 18 U/L,^[26] 30 U/L 52 to 57 U/L (20) 70 U/L.^[24,27] In normal gestation, ALT and AST are lower than non-pregnant age-matched women, however AST changes to lesser amount.^[23] Based on study of Cerutti *et al*, 1976 AST, ALT and GGT significantly increase during sixth month of pregnancy; however it is not obvious whether this is compared with early gestation.^[24] The primary fluctuations in liver function evaluation may be due to red cell demolition and ultimately happens liver injury.^[28] The results showed that liver damaged in pregnant women with preeclampsia. Although platelet count was nearly equal in both group, but

other biomarkers were higher in pregnant women with preeclampsia compared with normal pregnant women.

CONCLUSIONS

At the end of study, we conclude that pregnancy with preeclampsia likely results in HELLP syndrome. We suggest further studies to understand exact mechanism of problem.

Acknowledgments

We thank specially Young Researchers and Elite Club, Yasooj Branch, Islamic Azad University, Yasooj, Iran due to cooperation in this study.

Financial support and sponsorship

This study was supported by Young Researchers and Elite Club, Yasooj Branch, Islamic Azad University, Yasooj, Iran.

Conflicts of interest

There are no conflicts of interest.

References

1. Leung TN, Zhang J, Lau TK, Chan LY, Lo YD. Increased maternal plasma fetal DNA concentrations in women who eventually develop preeclampsia. *Clin Chem*. 2001;47(1):137-9.
2. Powe CE, Levine RJ, Karumanchi SA. Preeclampsia, a disease of the maternal endothelium the role of antiangiogenic factors and implications for later cardiovascular disease. *Circulation*. 2011; 123(24):2856-69.
3. Conde-Agudelo A, Belizan JM. Risk factors for pre-eclampsia in a large cohort of Latin American and Caribbean women. *BJOG*. 2000;107(1):75-83.
4. Leeman L, Fontaine P. Hypertensive disorders of pregnancy. *Am Fam Physician*. 2008;78(1). 93-100.
5. Sibai B, Dekker G, Kupferminc M. Pre-eclampsia. *The Lancet*. 2005;365(9461):785-99.
6. Levine RJ, Maynard SE, Qian C, Lim K-H, England LJ, Yu KF, et al. Circulating angiogenic factors and the risk of preeclampsia. *N Engl J Med*. 2004;350(7):672-83
7. Esplin MS, Fausett MB, Fraser A, Kerber R, Mineau G, Carrillo J, et al. Paternal and maternal components of the predisposition to preeclampsia. *N Engl J Med*. 2001;344(12):867-72.
8. Kashanian M, Baradaran HR, Bahasadri S, Alimohammadi R. Risk factors for pre-eclampsia: a study in Tehran, Iran. *Arch Iran Med*. 2011;14(6):412-5.
9. Riely CA. Liver disease in the pregnant patient. *Am J Gastroenterol*. 1999;94(7):1728-32.
10. Haram K, Svendsen E, Abildgaard U. The HELLP syndrome: clinical issues and management. A Review. *BMC pregnancy and childbirth*. 2009;9:8.
11. Arulkumaran N, Lightstone L. Severe pre-eclampsia and hypertensive crises. *Best Pract Res Clin Obstet Gynaecol*. 2013;27(6):877-84.
12. Weinstein L. Syndrome of hemolysis, elevated liver enzymes, and low platelet count: a severe consequence of hypertension in pregnancy. *Am J Obstet Gynecol*. 1982;142(2):159-67.
13. Abedi P, Mohaghegh Z, Afshary P, Latifi M. The relationship of serum vitamin D with pre-eclampsia in the Iranian women. *Matern Child Nutr*. 2014;10(2):206-12.
14. Shahbazian N, Shahbazian H, Ehsanpour A, Aref A, Gharibzadeh S. Hypertension and microalbuminuria 5 years after pregnancies complicated by pre-eclampsia. *Iran J Kidney Dis*. 2011;5(5):324-7.
15. Shirali S, Aghaei M, Shabani M, Fathi M, Sohrabi M, Moeinifard M. Adenosine induces cell cycle arrest and apoptosis via cyclinD1/Cdk4 and Bcl-2/Bax pathways in human ovarian cancer cell line OVCAR-3. *Tumour Biol*. 2013; 34(2):1085-95.
16. Barari A, Fakori Joybari M, Shirali S, Shojaee M, Khandandel A. Effect of eight-week consumption of purslane extract on peroxidase/antioxidant balance in women with type 2 diabetes. *MLJ*. 2014;8(2):1-7.
17. Sharifat M, Shirali S, Ebadi P. Investigation of changes in levels of AGEs and β 2-microglobulin in patients with type 2 diabetes. *JAAS*. 2014;4(9):1-10.
18. Mahdizadeh R, Shirali S, Ebadi P. Investigation of imbalance of trace elements in patients with type 2 diabetes mellitus. *JAAS*. 2014;4(9):11-21.
19. Borglin N. Serum transaminase activity in uncomplicated and complicated pregnancy and in newborns. *J Clin Endocrinol Metab*. 1958;18(8):872-7.
20. Romero R, Vizoso J, Emamian M, Duffy T, Riely C, Halford T, et al. Clinical Significance of Liver Dysfunction in Pregnancy-Induced Hypertension. *Obstetric Anesthesia Digest*. 1989;8(4):162.
21. Verhaeghe J, Anthony J, Davey D. Platelet count and liver function tests in proteinuric and chronic hypertension in pregnancy. *S Afr Med J*. 1991;79(10):590-4.
22. Hay JE. Liver disease in pregnancy. *Hepatology*. 2008;47(3):1067-76.
23. Shukla P, Sharma D, Mandal R. Serum lactate dehydrogenase in detecting liver damage associated with pre-eclampsia. *Br J Obstet Gynaecol*. 1978;85(1):40-2.
24. Cerutti R, Ferrari S, Grella P, Castelli G, Rizzotti P. Behaviour of serum enzymes in pregnancy. *Clin Exp Obstet Gynecol*. 1976;3:22-4.
25. Girling J, Dow E, Smith J. Liver function tests in pre-eclampsia: importance of comparison with a reference range derived for normal pregnancy. *Br J Obstet Gynaecol*. 1997;104(2):246-50.
26. VAN Dam PA, Renier M, Baekeland M, Buytaert P, Uyttenbroeck F. Disseminated intravascular coagulation and the syndrome of hemolysis, elevated liver enzymes, and low platelets in severe preeclampsia. *Obstet Gynecol*. 1989;73(1):97-102.
27. Goodlin R. Be ware the great imitator-severe preeclampsia. *Contemp Ob Gyn*. 1982;20:215.
28. McMahon L, O'Coighligh S, Redman C. Hepatic enzymes and the HELLP syndrome: a long-standing error? *Br J Obstet Gynaecol*. 1993;100(7):693-5.
